

OPIOID RISK TOOL (ORT)

		Mark each box that applies	Item score if FEMALE	Item score if MALE	
1	Family history of substance abuse	Alcohol	<input type="checkbox"/>	1	3
		Illegal drugs	<input type="checkbox"/>	2	3
		Prescription drugs	<input type="checkbox"/>	4	4
2	Personal history of substance abuse	Alcohol	<input type="checkbox"/>	3	3
		Illegal drugs	<input type="checkbox"/>	4	4
		Prescription drugs	<input type="checkbox"/>	5	5
3	Age (mark box if 16–45)	<input type="checkbox"/>	1	1	
4	History of preadolescent sexual abuse	<input type="checkbox"/>	3	0	
5	Psychological disease	Attention deficit disorder	<input type="checkbox"/>	2	2
		Obsessive compulsive disorder	<input type="checkbox"/>	2	2
		Bipolar	<input type="checkbox"/>	2	2
		Schizophrenia	<input type="checkbox"/>	2	2
		Depression	<input type="checkbox"/>	1	1
TOTAL			_____	_____	

Total Score Risk Category

0–3 = low risk

4–7 = moderate risk

≥8 = high risk

Reference: Webster LR. Predicting aberrant behaviors in opioid-treated patients: Preliminary validation of the opioid risk tool. *Pain Medicine*. 2005;6(6):432-442. Used with permission.

The ORT and other tools are available online at www.oregonpainguidance.org/clinical-tools.